



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Or, completing the multiplications,

$$(y^3 + z^3 - b^3)^3 + 27b^3y^3z^3 = 0.$$

Restoring the original values of  $y$ ,  $z$ , and  $b$ , we get,

$$27c(x^2 - a^2) = (2a - c)^3.$$

$$\text{Hence } x = \sqrt{\left(\frac{(2a - c)^3 + 27a^2c}{27c}\right)}.$$

This may be the root of the given equation or the root of any of the assumed equations, depending on the various values of  $a$  and  $c$ .

[This example is found in Bonycastle's Algebra (1845), page 97.]

Also solved by P. S. BERG, H. C. WILKES, and G. B. M. ZERR.

71. Proposed by F. P. MATZ, D. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

When  $x=0$ , find the limit of the expression

$$u = \left(\frac{m+x}{m-x}\right)^{\frac{1}{x}} + \left(\frac{m-x}{m+x}\right)^{\frac{1}{x}}.$$

I. Solution by O. W. ANTHONY, M. Sc., Columbian University, Washington, D. C., and G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas.

$$\text{Let } u = u_1 + u_2. \quad \therefore u_1 = \left(\frac{m+x}{m-x}\right)^{\frac{1}{x}},$$

$$\log u_1 = (1/x) \{ \log(m+x) - \log(m-x) \}$$

$$= (1/x) \{ [\log m + (x/m) - (x^2/2m^2) + (x^3/3m^3) - \dots] -$$

$$[\log m - (x/m) - (x^2/2m^2) - (x^3/3m^3) - \dots] \}$$

$$= \frac{2}{x} \left( \frac{x}{m} + \frac{x^3}{3m^3} + \frac{x^5}{5m^5} + \dots \right) = 2 \left( \frac{1}{m} + \frac{x^2}{3m^3} + \frac{x^4}{5m^5} + \dots \right)$$

$$= 2/m, \text{ when } x=0. \quad \log u_2 = -\log u_1 = -2/m, \text{ when } x=0.$$

$$\therefore u_1 = e^{2/m}, u_2 = e^{-2/m}. \quad \therefore u = e^{2/m} + e^{-2/m} \text{ when } x=0.$$

II. Solution by H. C. WHITAKER, M. Sc., Ph. D., Professor of Mathematics in Philadelphia Manual Training School, Philadelphia, Pennsylvania.

Since  $(1+x)^{1/x} = e$  when  $x=0$ , we have

$$\left(\frac{m+x}{m-x}\right)^{\frac{1}{x}} = \left[ \left(1 + \frac{2x}{m-x}\right)^{\frac{m-x}{2x}} \right]^{\frac{2}{m-x}} = e^{2/m} \text{ when } x=0.$$

In the same way  $\left(\frac{m-x}{m+x}\right)^{\frac{1}{x}} = e^{-2/m}$  when  $x=0$ . Hence  $u = e^{2/m} + e^{-2/m}$ .

Also solved by J. SCHEFFER.

## GEOMETRY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

### SOLUTIONS OF PROBLEMS.

69. Proposed by WILLIAM SYMMONDS, M. A., Professor of Mathematics and Astronomy in Pacific College, Santa Rosa, California; P. O., Sebastopol, California.

To divide a square card into right-lined sections in a manner, that a rectangle of a given breadth can be formed from the sections; likewise, form a square from a rectangular card.

I. Solution by G. B. M. ZERR, A. M., Ph. D., Texarkana, Arkansas.

(1). Let  $ABCD$  be the square. Produce  $DA$  to  $H$  making  $AH$  equal the given width of the rectangle, join  $HB$ , and draw  $KO$  perpendicular to  $HB$  at its mid-point, then  $O$  is the center of the circle through  $HB$ . Produce  $AD$  to meet circle at  $G$ ;  $AG$  is the length of the required rectangle. Take  $AE=AH$  and complete the rectangle  $AEFG$ .

Now the right triangle

$AHB$  = right triangle  $BCN$  = right triangle  $MFG$ .


$\therefore CN=AE$  and  $DN=BE$ ;

$\therefore \triangle BEM = \triangle DNG$ .

$\therefore ABCD = ADNME + BCN + BEM$

$= ADNME + MFG + NDG = AEFG$ .

(2). Let  $AEFG$  be the given rectangle. Produce  $GA$  to  $H$  making  $AH=AE$ . Upon  $HG$  describe the semi-circle. Then  $AB$  is a side of the required square. Complete the square  $ABCD$  and draw  $BG$ . The rest of the proof is the same as above.



II. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.

(1). Let  $ABCD$  represent the square card. From  $A$  lay off on  $AB$  the width of the rectangle successively as many times as possible, as  $AE$ ,  $EF$ .

Then from the opposite corner  $C$ , lay off *one width only* of the rectangle, as  $CG$ . Now cut through on line  $GB$ . Then cut  $FH$  and  $EI$  parallel to  $DA$ .

